

Bryum caespitium

Almost cosmopolitan.

Bryum campylohectum

Australia, New Zealand.

Campylopus introflexus

Australia, New Zealand, Pacific Islands, introduced to UK, N. & S.

America, S. Africa.

Weisia controversa

Cosmopolitan.

Cephalozia arctica

Southern Australia, New Zealand.

There is some confusion over the use of *Bryum albo-limbatum* in Australian literature. Elsewhere it is recognised as a synonym for *Bryum capillare*, which this Australian moss is not. It is possible that it is in fact, *Bryum andicola*, if so the distribution would be: N. S. and central America, E. and S. Africa and southern Australia. Further collection of this species may provide information that could clarify the above riddle.

CONCLUSIONS

This bryophyte flora list although not considered comprehensive, when used in conjunction with other articles produced on this reserve

(Turpin, 1991, Cranfield & Parker, 1992) will enable the readers to develop an overall understanding of the natural dynamics involved.

ACKNOWLEDGEMENTS

We wish to thank Dr N.G. Marchant for his guidance.

REFERENCES

- Bonner, C.E.B. 1977. *Index Hepaticarum*. Pars VIIa: Supplement A-C.
Catcherside, D.G. 1980. *Mosses of South Australia*. Woolman, Govt. Printer South Australia.
Cranfield, R.J. & Parker, C.M. 1992. Flowering Calendar for Reserve No. 3694 in Metropolitan Perth. *West. Aust. Nat.* 19: 48-60.
Magill, R.E., 1981, 1987. *Flora of Southern Africa*. Part 1, Fascicles 1 & 2. Botanical Research Institute, Dept. of Agriculture and Fisheries, Republic of South Africa.
Turpin, M.C. 1991. Ecological appraisal of an isolated Banksia Woodland, Reserve No. 3694, south of the Swan River, Perth. *West. Aust. Nat.* 18: 131-138.

DIET OF VARANUS CAUDOLINEATUS (REPTILIA: VARANIDAE)

By GRAHAM G. THOMPSON

Edith Cowan University, Joondalup, 6027, Western Australia, and Zoology Department, University of Western Australia, Nedlands, 6009, Western Australia

and DENNIS KING

c/- Western Australian Museum, Francis St, Perth, 6000, Western Australia

ABSTRACT

The diet of museum specimens of *Varanus caudolineatus* is significantly different from those caught on Atley Station in Western Australia. The museum specimens had a much broader range of prey types, (predominantly spiders, grasshoppers and lizards) than those found on Atley Station that had a very high predominance of scorpions and a lower, but significant number of terrestrial spiders in their stomachs and intestines. These data suggest that *V. caudolineatus* at the Atley site essentially forage primarily on the ground rather than in trees as might be expected for an arboreal lizard.

INTRODUCTION

Varanids are generally reported as being carnivorous and opportunistic feeders, eating predominantly invertebrates. However, a few species specialise on prey types that reflect their habitat (Losos and Greene 1988). Diets of many species of varanids have been determined from museum specimens (Pianka 1968, 1969, 1970 a, b, 1971, Greene 1986, Losos and Greene 1988, James *et al.* 1992) or from freshly caught specimens from a diverse range of locations (King and Green 1979, Pianka 1968, 1969, 1970a 1971, 1986).

There are few reports of the diet of varanids based on stomach contents collected from a single location

(Shine 1986, Thompson in press). Significant ecological information is often inferred from dietary data.

Varanus caudolineatus is an arboreal pygmy goanna that usually takes refuge under loose bark or in the hollows of mulga trees; it is not usually found in spinifex or sand plain habitats (Pianka 1969). Its diet is quite varied and includes mainly grasshoppers, roaches and geckos.

This study reports the stomach contents of 88 Western Australian Museum (WAM) specimens of *V. caudolineatus* and contrasts this data with diet of 33 specimens caught in December 1991 in close proximity on Atley Station, W.A. (119°07'E, 28°25'S), 4 caught in July 1991 North

of Ajana, W.A. (114°45'E, 27°35'S) and the 78 other specimens (largely from the W.A. and S.A. Museum) previously examined by Pianka (1969).

METHODS AND MATERIALS

A ventral incision was made in museum specimens (W/AM) and the gut contents removed and identified. Stomach contents or scats were collected from 24 individual *V. caudolineatus* caught in tree hollows and pit traps on Atley Station and four *V. caudolineatus* caught in tree hollows in the area north of Ajana. Nine of the monitors were subsequently recaptured at least five days later on Atley Station and their stomach contents or scats were again collected. Stomach contents were flushed with tepid water, using a technique similar to that reported by Legler and Sullivan (1979). Scats were collected from the calico bags used to temporarily hold *V. caudolineatus* on the Atley and Ajana sites. The stomach contents and scats were stored in a dilute solution of formalin (approximately 10%) for later identification. Neither stomach contents nor scats were collected from seven of the 33 lizards examined from the Atley site. Three of the four lizards from Ajana provided prey items from their stomach.

The Atley Station site contains red loamy soils which supports a vegetation consisting mainly of spinifex (*Triodia* sp.), grasses, mulga (*Acacia* sp.) and gum (*Eucalyptus* sp.) trees. A representative sample of the potential small vertebrate and invertebrate prey items were collected from 60 pit traps (20 litre plastic containers) with 10m drift

fences installed on the Atley site.

The analysis of the difference between the stomach contents of museum specimens and those from the Atley site was done using the SPSS/PC discriminant analysis program.

RESULTS

The stomachs of 46 of 88 museum specimens examined were empty. The contents of the remaining 42 stomachs contained predominantly spiders, grasshoppers and lizards (Table 1). Seven of the lizards were identified as geckos, *Gehyra* sp., and one stomach contained at least two *Gehyra* sp. tails. Most stomachs contained only one item but four contained two or more identifiable items.

Three stomachs of the 4 *V. caudolineatus* from the site north of Ajana contained prey items; these were a spider, a beetle and a cricket (Table 1).

V. caudolineatus from the Atley site fed predominantly on scorpions, followed by a lesser number of large spiders, that were often found in holes in the ground during the day, and lizards (Table 1). Two of the lizards were skinks, one was a juvenile *Menidia greyii* the other was the dismembered parts of a small *Ctenopus schomburgkii*. In addition, three *V. caudolineatus* from the Atley site and one from the site north of Ajana had red gravel in their stomachs. This was probably ingested while they were capturing prey on the ground.

There is a significant difference (chi-square 74.63, df 6, $p < 0.0001$) in the gut contents of the museum

Table 1. The number of stomachs (and scats*) with the incidence of prey items in *V. caudolineatus*

	Museum specimens examined	Pianka 1969	Atley site*	Ajana site
No specimens examined	88	78	33	4
No stomachs empty	46	44	7	1
PREY TYPE				
Centipedes	1	2	1	
Scorpions	10	6	17	
Spiders	8	10	7	1
Grasshoppers	3	11	2	
Roaches		1	1	
Moths	4	4	0	
Larva	1		1	
Beetles	8	6	0	1
Unidentified invertebrate	1			
Egg sac				
Bee		1		
Cricket		1		
Lizards - egg sac		1		
- other parts	9	10	3	
Twigs		2	1	
Red gravel		3	1	

* the contents of lizard scats are included in the data from the Atley site.

specimens we examined and those caught on the Atley Station. 91.2% of the gut contents were classified correctly with the group centroids being 1.95 for the Atley site and -1.13 for the museum specimens.

Potential prey items captured in the 60 pit traps located on the Atley site were placed into three groups, depending on their relative abundance in the traps (Table 2). In addition, *Gehyra variegata* and *Egernia depressa* were found in relatively low number in trees in which *V. caudolineatus* were found.

DISCUSSION

There is a similarity between the stomach contents of the museum

specimens we examined and those examined by Pianka (1969) with the exception that Pianka's sample contained a relatively higher number of roaches. This similarity would have been expected.

There is however a significant difference between the diets of *V. caudolineatus* found on the Atley site, and those examined by us in the Western Australian Museum collection. The very high predominance of scorpions and lower, but significant number of terrestrial spiders in the stomachs and intestines of *V. caudolineatus* from the Atley site contrasts with the much broader range of prey types in the museum specimens we examined and in Pianka's (1969)

Table 2. Relative abundance of potential prey items found in pit traps on the Atley site.

Item	Pit trap catch frequency		
	Often caught	Seldom caught	Rarely caught
INSECTA			
Thysanura (silverfish)		*	
Blattodea (roaches)		*	
Isopoda (termites)			*
Mantodea (preying mantids)			*
Orthoptera (grasshoppers)		*	
Phasmatodea (stick insects)			*
Coleoptera (beetles)	*		
Hymenoptera (ants)	*		
ARACHNIDA			
Scorpionida (scorpions)	*		
Araneae (spiders)	*		
CHILOPODA			
Scolopendrida (centipedes)	*		
REPTILIA			
Scincidae (skinks)	*		
Gekkonidae (geckos)		*	
Varanidae (goanna)		*	

samples. In this particular circumstance the stomach contents of specimens coming from a variety of locations [museum and Pianka (1969)] do not necessarily provide a good indication of the diet of the same species at a particular site. This is similar to the situation for *V. merriami* (Shine 1986).

The different diet of the small monitors at the Atley site probably reflects the food items available and able to be captured and ingested. Centipedes were abundant there, but the adults may be too large or too difficult to subdue. Many of the beetles present there might also have been too large and their exoskeletons too difficult to penetrate

for them to become prey for *V. caudolineatus*. The small skinks were caught in the pit-traps were predominantly *Ctenotus schomburkii* and *Ctenotus leonhardi*, while the predominant gecko was *Rhynchoedura ornata*. *C. schomburkii* and *R. ornata* appear to be within the suitable prey size range for *V. caudolineatus* to ingest but may be too difficult to locate and capture. *C. leonhardi* is a larger skink and may be too fast and too large for *V. caudolineatus* to capture. *Gehyra variegata*, which was relatively abundant in the stomachs of museum specimens, but was not found in the stomachs of the Atley site monitors, was caught only once

in the pit-traps, but were occasionally found under the bark of trees.

All identifiable species of scorpions and spiders found in the stomachs and scats of these small monitors are ground dwelling invertebrates. In addition, both *M. greyii* and *C. schomburkii* are both ground dwelling skinks that live around the base of spinifex clumps or in holes in the ground. This supports other behavioural data (Thompson, unpublished) that indicate that *V. caudolineatus* retreat to trees when threatened, to sleep or to observe their surrounds, but essentially forage on the ground.

No *V. caudolineatus* were captured at night in the pit-trapping program on the Atley site, although the ambient temperature at night was often in the mid-to-high 30°Cs. Scorpions, and many of the spider species that were preyed-upon, were only found out of their burrows after dark. On two occasions in soft red loamy soils clear tracks of *V. caudolineatus* were found leading to the entrance of scorpion holes, suggesting that these lizards enter burrows in search of prey. The red gravel in the stomachs of monitors may be ingested when they are eating scorpions or spiders captured in burrows.

This study provides further support for the point made by Shine (1986) and Thompson (in press) that the diets of varanids vary significantly with locality and time of the year. Much of the dietary data and the derived ecological information for varanids comes from museum specimens (Losos 1988, Greene 1986, James *et al.* 1992). It would therefore appear inappropriate to draw

specific conclusions about the diets of varanids at a particular location from museum specimens or from those caught in other locations. It would be interesting to know however, if there was a relationship between the density of *V. caudolineatus* and the high availability of scorpions at the Atley site or whether this small monitor would feed on other invertebrates in the absence of scorpions.

Contrary to Pianka's (1969) suggestion that *V. caudolineatus* are not found in spinifex habitats, most of the Atley sites from which *V. caudolineatus* were collected had patches of spinifex ground cover and a sparse cover of mulga trees.

ACKNOWLEDGEMENTS

The field work component of this research was made possible by the financial assistance of Edith Cowan University and the University of Western Australia. The assistance of P. Harely, W. Thompson and S. Thompson in digging in pit-traps, catching *V. caudolineatus* and checking pit-traps was very much appreciated. The comments of P. C. Withers significantly improved the manuscript. K. Applin and L. Smith provided access to the collections of the Western Australian Museum. The approval by R. and J. Broadhurst to have access to Atley Station was appreciated. All lizards were caught under licence issued by the Department of Conservation and Land Management to G. Thompson. Animal experimentation on live specimens was done with the approval of the Animal Welfare Committee of the University of Western Australia.

REFERENCES

- GREENE, H.W., 1986. Diet and Arboreality in the Emerald Monitor, *Varanus prasinus*, with Comments on the Study Adaptation. *Fieldiana Series No 31*: i-12.
- KING, D. and GREEN, B. 1979. Notes on Diet and Reproduction of the Sand Goanna, *Varanus gouldii rosenbergi*, *Copeia*, 1979: 64-70.
- JAMES, C.D., LOSOS, J.B. and KING, D.R. 1992. Reproductive biology and diets of goannas (Reptilia: Varanidae) from Australia. *J. Herpetology*, 26: 128-136.
- LEGLER, J.M. and SULLIVAN, L.J. 1979. The Application of Stomach-Flushing to Lizards and Anurans, *Herpetologica*, 35: 107-110.
- LOSOS, J.B. and GREENE, H.W. 1988. Ecological and evolutionary implications of diet in monitor lizards, *Brit. J. Linnean Soc.*, 35: 379-407.
- PIANKA, E.R. 1968. Notes on the Biology of *Varanus eremius*, *West. Aust. Nat.*, 11: 39-44.
- PIANKA, E.R. 1969. Notes on the Biology of *Varanus caudolineatus* and *Varanus gilleani*, *West. Aust. Nat.*, 11: 76-82.
- PIANKA, E.R. 1970a. Notes on the biology of *Varanus gouldii flaviventris*, *West. Aust. Nat.*, 11: 141-144.
- PIANKA, E.R. 1970b. Notes on *Varanus brevicauda*, *West. Aust. Nat.*, 11: 113-116.
- PIANKA, E.R. 1971. Notes on the biology of *Varanus tristis*, *West. Aust. Nat.*, 11: 180-183.
- PIANKA, E.R. 1986. *Ecology and Natural History of Desert Lizards*, Princeton Uni. Press, Princeton, NJ.
- SHINE, R. 1986. Food habits, habitat and reproductive biology of four sympatric species of varanid lizards in tropical Australia, *Herpetologica*, 42: 346-360.
- THOMPSON, G.G. (in press). Notes on the diet of *Varanus gouldii* in a semi urban environment, *West. Aust. Nat.*

D.L. SERVenty'S GOOSEBERRY HILL PLANT LIST

By B. M. J. HUSSEY

Department of Conservation and Land Management, PO Box 104,
Como, Western Australia 6152.

INTRODUCTION

The late D. L. Serventy worked at the CSIRO laboratory in Helena Valley, and owned Lot 306, The Knoll, Gooseberry Hill, which he used as a weekend cottage. At his behest, G. M. Storr of the WA Museum collected 74 plants on The Knoll on 5th October 1957. Dr Serventy continued sporadic collections at the locality until 1967, reaching a total of 92 taxa. The specimens collected by Storr were identified by C. A. Gardner, Serventy's by R.D. Royce or A.S. George, all of the WA Herbarium.

I became aware of the existence of the collection when Dr Serventy permitted me to use the house as a base for school biology/geography excursions during the 1970s. After his death, the Serventy family kindly gave me the collection to assist with my study of the flora of Gooseberry Hill. It is described here as a distinct historical collection, and the specimens have been lodged in the WA Herbarium [PERTH].

A list of the plants represented in the collection is given in Table 1. The arrangement and nomenclature follows Marchant *et al.* (1987) but where taxonomic changes have occurred, the name originally applied is given in brackets.

DISCUSSION

Gooseberry Hill is a promontary of lateritic caprock which forms the southern edge of the Helena Valley where it cuts through the Darling Scarp. The site, geology, soil and vegetation has been described in Hussey (1993). Basically, it is a narrow plateau of massive laterite, breaking sharply away on the north, west and south sides to form steep scree slopes. The vegetation is open forest of Jarrah, *Eucalyptus marginata*, and Marri, *Eucalyptus calophylla*, with a heath understorey.

D. L. Serventy's block, occupied part of the caprock and scree slope, facing north over the Helena Valley and the CSIRO laboratories where he worked. At the time when the collection was made very few blocks were developed, now almost all are. With the exception of two species, *Hovea pungens* and *Conostylis androstemma*, the specimens come from the laterite area, possibly even from Lot 306 itself.

Currently, the species list for the lateritic section of the public land on The Knoll stands at 188 species. Since this collection contains only 90 plants, it is clearly not exhaustive - there are only three members of the Proteaceae, for example, and no Myrtaceae at all. It cannot therefore